bs-11456R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Dscam Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 1826 **SWISS:** 060469

Target: Dscam

Immunogen: KLH conjugated synthetic peptide derived from human Dscam:

301-400/2012. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: DSCAM is a cell adhesion molecule belonging to the

immunoglobulin superfamily. It is predominantly expressed in brain and contains six Fibronectin type-III domains and ten Ig-like C2-type domains. In mice, DSCAM is responsible for regulating isoneuronal self-avoidance, the tendency for sister arbors to avoid crossing each other and to spread out proportionately over an area. Isoneuronal self-avoidance is important for proper terminal branching (arborization). In some cell types, DSCAM also mediates heteroneuronal self-avoidance, which is important for the regular spacing of cell bodies and the prevention of hyperfasciculation. In humans, two DSCAM isoforms exist due to alternative splicing. The long isoform (also known as CHD2-42) is a single pass type I membrane protein, while the short isoform (CHD2-52), which lacks the C-terminal transmembrane containing region (amino acids 1572-2012), is secreted.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500)


Reactivity: Mouse, Rat

(predicted: Human, Sheep, Cow, Chicken, Dog, Horse)


Predicted MW.: 220 kDa

Subcellular Secreted ,Cell membrane

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Dscam) Polyclonal Antibody, Unconjugated (bs-11456R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Dscam) Polyclonal Antibody, Unconjugated (bs-11456R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.