bs-15489R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

HIGD1B Rabbit pAb

GenelD: 51751 **SWISS:** Q9P298

Target: HIGD1B

Immunogen: KLH conjugated synthetic peptide derived from human HIGD1B:

22-80/99.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a member of the hypoxia inducible gene 1

(HIG1) domain family. The encoded protein is localized to the cell membrane and has been linked to tumorigenesis and the progression of pituitary adenomas. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2012]

Applications: WB (1:500-2000)

400-901-9800

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)

Reactivity: (predicted: Human, Mouse,

Rat, Rabbit, Sheep, Cow,

Dog, Horse)

Predicted MW.: 11 kDa

Subcellular Location: Cell membrane

- SELECTED CITATIONS -

• [IF=3.5] Xiangyu Chen. et al. Downregulation of HIGD1B induces mitochondria-mediated apoptosis in gastric cancer cells by inactivating Akt and ERK pathways. DRUG DEVELOP RES. 2024 Jun;85(4):e22221 WB,IHC; Human. 38863387