## bs-23116R

## [ Primary Antibody ]

## NR2E1 Rabbit pAb



		400-901-9800
Host: Rabbit	lsotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal	GenelD: 7101	
Target: NR2E1		
Immunogen: KLH conjugated syn 151-250/451.	thetic peptide derived from human NR2E1:	
Purification: affinity purified by F	rotein A	
<ul> <li>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</li> <li>Background: NR2 proteins are a large family of nuclear hormone receptor transcription factors. The proteins belonging to this family are characterized by discrete domains functioning in DNA and ligand binding. NR2E1 (nuclear receptor subfamily 2, group E, member 1), also known as TLX, is an essential component in the formation of synaptic plasticity and dendritic structure in retinal astrocytes. In addition, NR2E1 is a orphan receptor that binds DNA as part of the hormone response element (HRE), a transcription regulator for hormones. DNA-binding orphan receptors have the conserved sequence 5'-AAGGTCA-3', a motif that determines substrate binding specificity. NR2E1 is expressed in brain tissue, with highest levels in astrocytes, and is localized to the nucleus. Mutations in the gene that encodes NR2E1 may lead to retinal dystrophy, a disorder characterized was reduction in the thickness of the retina</li> </ul>		Reactivity: Mouse, Rat (predicted: Human, Rabbit, Pig, Cow, Dog) Predicted 43 kDa MW: 43 kDa Subcellular Location: Nucleus e st

## - VALIDATION IMAGES ------



Sample: Brain (Mouse) Lysate at 40 ug Brain (Rat) Lysate at 40 ug Primary: Anti-NR2E1 (bs-23116R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 43 kD Observed band size: 43 kD