

bs-20761R**[Primary Antibody]****HTRA2 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse (predicted: Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse) Predicted MW.: 35 kDa Subcellular Location: Cell membrane ,Cytoplasm
Clonality: Polyclonal		
GeneID: 27429	SWISS: O43464	
Target: HTRA2		
Immunogen: KLH conjugated synthetic peptide derived from human HTRA2: 331-430/458.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene encodes a serine protease. The protein has been localized in the endoplasmic reticulum and interacts with an alternatively spliced form of mitogen-activated protein kinase 14. The protein has also been localized to the mitochondria with release to the cytosol following apoptotic stimulus. The protein is thought to induce apoptosis by binding the apoptosis inhibitory protein baculoviral IAP repeat-containing 4. Nuclear localization of this protein has also been observed. Alternate splicing of this gene results in two transcript variants encoding different isoforms. Additional transcript variants have been described, but their full-length sequences have not been determined. [provided by RefSeq, Jul 2008]		

— VALIDATION IMAGES —

Sample: Lovo Cell Lysate at 40 ug Jurkat Cell Lysate at 40 ug
Primary: Anti- HTRA2 (bs-20761R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 35kD
Observed band size: 35 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain);
Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min;
Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes;
Blocking buffer (normal goat serum) at 37°C for 30min;
Antibody incubation with (HTRA2) Polyclonal Antibody, Unconjugated (bs-20761R) at 1:400 overnight at 4°C,
followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain glioma);
Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min;
Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes;
Blocking buffer (normal goat serum) at 37°C for 30min;
Antibody incubation with (HTRA2) Polyclonal Antibody, Unconjugated (bs-20761R) at 1:400 overnight at 4°C,
followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.