

**bs-20763R****[ Primary Antibody ]****PARP1 Rabbit pAb**

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**DATASHEET****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 142**SWISS:** P09874**Target:** PARP1**Immunogen:** KLH conjugated synthetic peptide derived from human PARP1: 581-650/1014.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** This gene encodes a chromatin-associated enzyme, poly(ADP-riboseyl)transferase, which modifies various nuclear proteins by poly(ADP-riboseyl)ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transformation and also in the regulation of the molecular events involved in the recovery of cell from DNA damage. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pathophysiology of type I diabetes. [provided by RefSeq, Jul 2008].

**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (1ug/Test)


**Reactivity:** Human, Mouse, Rat  
(predicted: Rabbit, Pig, Sheep, Cow)

**Predicted MW.:** 112 kDa


**Subcellular Location:** Nucleus

**VALIDATION IMAGES**


Sample: Testis(Mouse) Lysate at 40 ug Primary: Anti-PARP1 (bs-20763R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 112 kD  
Observed band size: 112 kD



Sample: Lane 1: K562 (Human) Cell Lysate at 30 ug Lane 2: SH-SY5Y (Human) Cell Lysate at 30 ug Lane 3: U251 (Human) Cell Lysate at 30 ug Lane 4: HeLa (Human) Cell Lysate at 30 ug Primary: Anti-PARP1 (bs-20763R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 115 kD  
Observed band size: 115 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PARP1) Polyclonal Antibody, Unconjugated (bs-20763R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control:293T. Primary Antibody (green line): Rabbit Anti-PARP1 antibody (bs-20763R)  
Dilution: 1µg /10<sup>6</sup> cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution:

**Important Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

---

## — SELECTED CITATIONS —

- **[IF=3.509]** Xin Shan. et al. Proteomic analysis of healthy and atretic porcine follicular granulosa cells. J Proteomics. 2021 Feb;232:104027 WB ;Pig. 33130110
- **[IF=3.8]** Cuicui Zhuang. et al. Escherichia coli infection mediates pyroptosis via activating p53-p21 pathway-regulated apoptosis and cell cycle arrest in bovine mammary epithelial cells. MICROB PATHOGENESIS. 2023 Sep;:106338 WB ;Bovine. 37683833