

bs-9194R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

IGSF1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200) Reactivity: Mouse (predicted: Human, Rat, Pig, Dog, Horse) Predicted MW.: 80, 144 kDa Subcellular Location: Secreted ,Cell membrane
Clonality: Polyclonal		
GeneID: 3547	SWISS: Q8N6C5	
Target: IGSF1		
Immunogen: KLH conjugated synthetic peptide derived from human IGSF1: 271-380/1336. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: IGSF1 is a highly glycosylated immunoglobulin domain-containing protein. IGSF1 has been shown to act as a coreceptor in inhibin signaling, however, it does not appear to be a high-affinity inhibin receptor by itself. May reduce or inhibit activin A signaling and is believed to be necessary in the mediation of specific effects of inhibin B on activin-stimulated transcription. IGSF1 has been found to interact with several members of the ACVR family and possibly some members of the BMPR group. There are 3 known isoforms of IGSF1, with 1 and 2 likely being multi-pass membrane proteins. Isoform 3 is believed to be expressed as a secreted form. Expression is high in pancreas, testis and fetal liver, while heart, prostate and small intestine show only moderate expression. IGSF1 may be found at very low levels in brain, muscle, thymus, ovary, colon, fetal lung and fetal kidney. Isoform 3 has been detected in pituitary gland.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (IGSF1) Polyclonal Antibody, Unconjugated (bs-9194R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.