

bs-0295D-Cy3

[Secondary Antibodies]

Donkey Anti-Rabbit IgG H&L, Cy3 conjugated

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

<p>Host: Donkey</p> <p>Clonality: Polyclonal</p> <p>Target: Donkey Anti-Rabbit IgG H&L</p> <p>Purification: affinity purified by Protein G</p> <p>Concentration: 2.0 mg/ml</p> <p>Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.</p>	<p>Isotype: IgG</p> <p>Applications: IF (1:100-1000) Excitation Spectrum: 552nm Emission spectrum: 570nm</p> <p>Reactivity: Rabbit</p>
--	---

— SELECTED CITATIONS —

- **[IF=6.388]** Liang Liang. et al. Oridonin relieves depressive-like behaviors by inhibiting neuroinflammation and autophagy impairment in rats subjected to chronic unpredictable mild stress. PHYTOTHER RES. 2022 Jun 09 IF ;Rat. 35686337
- **[IF=5.1]** Zhu Yuan-Zheng. et al. Genome-wide search links senescence-associated secretory proteins with susceptibility for coronary artery disease in mouse and human. J GERONTOL A-BIOL. 2024 Feb;; IF ;Mouse. 38416803
- **[IF=4.2]** Ruixiao Guo. et al. The GABAergic pathway from anterior cingulate cortex to lateral hypothalamus area regulates irritable bowel syndrome in mice and its underlying mechanism. J NEUROCHEM. 2024 Jun;; IF ;Mouse. 38877776