
KAZALD1 Rabbit pAb

Catalog Number: bs-17082R

Target Protein: KAZALD1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Rat (predicted: Mouse, Rabbit, Pig, Cow, Dog, Horse)

Predicted MW: 30 kDa

Entrez Gene: 81621

Swiss Prot: Q96I82

Source: KLH conjugated synthetic peptide derived from human KAZALD1: 71-170/304.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: BONO1, also known as KAZALD1 (kazal-type serine protease inhibitor domain-containing protein 1), IGFBP-rP10, FKSG28 or FKSG40, is a 304 amino acid secreted extracellular matrix protein that promotes matrix assembly. BONO1 is expressed in developing bones and odontoblasts in teeth, where it plays a role in osteoblast proliferation during bone formation and regeneration. BONO1 is also expressed at high levels in spleen, and is found at lower levels in lung, skin, urinary bladder, brain, tongue, kidney and large intestine. Existing as two alternatively spliced isoforms, BONO1 contains one Kazal-like domain, an IGFBP N-terminal domain and a single Ig-like C2-type (immunoglobulin-like) domain. The gene encoding BONO1 maps to human chromosome 10, which contains over 800 genes and 135 million nucleotides. Cockayne syndrome, Cockayne syndrome and trisomy 10 are associated with defects in chromosome 10.


VALIDATION IMAGES



Sample: DU145(Human) Cell Lysate at 40 ug Primary: Anti-KAZALD1 (bs-17082R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 30 kD Observed band size: 30 kD



Paraformaldehyde-fixed, paraffin embedded (Rat small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KAZALD1) Polyclonal Antibody, Unconjugated (bs-17082R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KAZALD1) Polyclonal Antibody, Unconjugated (bs-17082R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.