## bs-0295D-BF647

# [ Secondary Antibodies ]

# Donkey Anti-Rabbit IgG H&L, BF647 conjugated

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET

Host: Donkey Isotype: IgG

Clonality: Polyclonal

Target: Donkey Anti-Rabbit IgG H&L Purification: affinity purified by Protein G

Concentration: 2.0 mg/ml

**Storage:** 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50%

glycerol.

Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.

Applications: IF (1:100-1000)

Flow-Cyt (1:100-1000) ICC/IF (1:100-1000)

Excitation Spectrum: 647nm Emission spectrum: 666nm

Reactivity: Rabbit

### - SELECTED CITATIONS -

• [IF=17] Manon Rivagorda. et al. A primary cilia-autophagy axis in hippocampal neurons is essential to maintain cognitive resilience.NATURE AGING.2025 Feb 21. IF; Rabbit. 39984747