bs-12588R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

PDE8B Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 8622 SWISS: 095263

Target: PDE8B

Immunogen: KLH conjugated synthetic peptide derived from human PDE8B:

351-450/885.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a cyclic nucleotide

phosphodiesterase (PDE) that catalyzes the hydrolysis of the second messenger cAMP. The encoded protein, which does not hydrolyze cGMP, is resistant to several PDE inhibitors. Defects in this gene are a cause of autosomal dominant striatal degeneration (ADSD). Several transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2010]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) ICC/IF (1:100-500) **ELISA** (1:5000-10000)

Reactivity: (predicted: Human, Mouse,

Predicted MW.: 99 kDa

Subcellular Location: Cytoplasm

- SELECTED CITATIONS -

• [IF=4.4] Qiu Nian-zhuang. et al. Phosphodiesterase 8 (PDE8): Distribution and Cellular Expression and Association with Alzheimer's Disease. NEUROCHEM RES. 2024 May;:1-12 IF; Mouse. 38782837