

**bsm-1240M****[ Primary Antibody ]****human Fibrinogen Mouse mAb****BioSS**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —****Host:** Mouse**Isotype:** IgG1**Clonality:** Monoclonal**CloneNo.:** 4F7**GeneID:** 2243**SWISS:** P02671**Target:** human Fibrinogen**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** Size : 50ul/100ul/200ul


0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Size : 200ug (PBS only)

0.01M PBS

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** Fibrinogen is the main protein of blood coagulation system. It is a large protein and it consists of two identical subunits that contain three polypeptide chains: alpha, beta and gamma. All chains are connected with each other by a number of disulfide bonds. Fibrinopeptides A (1 to 16 amino acids) and B (1 to 17 amino acids) are released by thrombin from the N terminal parts of alpha and beta chains, respectively. In this way fibrinogen is converted into fibrin, which by means of polymerization forms a fibrin clot. Fibrinogen clotting underlies pathogenesis of MI, thromboembolism and thromboses of arteries and veins, since fibrin is the main substrate for thrombus formation. Fibrinogen activation is also involved in pathogenesis of inflammation, tumor growth and many other diseases. The normal fibrinogen concentration in plasma is about 3 mg/ml. The elevated level of fibrinogen in patient's blood is regarded as an independent risk factor for cardiovascular diseases. An increase in blood fibrinogen concentration was shown to be a strong predictor of coronary heart disease (Sonel A. et al, and Rapold H.J. et al). All these facts make fibrinogen an important parameter in the diagnosis of cardiovascular diseases.

**Applications:** Flow-Cyt (1ug/Test)**Reactivity:** Human**Predicted MW.:** 340 kDa**Subcellular Location:** Secreted**— VALIDATION IMAGES —**

Blank control (black line) :HepG2. Primary

Antibody (green line): Mouse Anti-human

Fibrinogen antibody (bsm-1240M)

Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-mouse IgG-AF488 Dilution:

0.5ug/Test. Isotype control (orange line) :

Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then

incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary

Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room

temperature. Acquisition of 20,000 events was

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

performed.