

bs-0295M-FITC**[Secondary Antibodies]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Mouse Anti-Rabbit IgG H&L, FITC conjugated**— DATASHEET —**

Host: Mouse Clonality: Polyclonal Target: Mouse Anti-Rabbit IgG H&L Purification: affinity purified by Protein G Concentration: 2.0 mg/ml Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 10 ¹¹ variants.	Isotype: IgG Applications: IF (1:100-1000) Flow-Cyt (1:100-1000) ICC/IF (1:100-1000) Excitation Spectrum: 495nm Emission spectrum: 519nm Reactivity: Rabbit
--	--

— SELECTED CITATIONS —

- **[IF=8.897]** Wu, Qingsi. et al. Targeted delivery of celastrol to glomerular endothelium and podocytes for chronic kidney disease treatment. Nano Res. 2021 Dec;;1-13 IF ;Mouse. 34925707
- **[IF=8.3]** Ling Guo. et al. Glomerulus-Targeted ROS-Responsive Polymeric Nanoparticles for Effective Membranous Nephropathy Therapy. ACS APPL MATER INTER. 2024;XXXX(XXX):XXX-XXX IF ;Mouse. 38940537
- **[IF=6.208]** Dezhang Lu. et al. The Mechanism of Lipopolysaccharide's Effect on Secretion of Endometrial Mucins in Female Mice during Pregnancy. INT J MOL SCI. 2022 Jan;23(17):9972 IHC ;Mouse. 36077364
- **[IF=3.12]** Dong, Feng, et al. "The isolation and characterization of a telomerase immortalized goat trophoblast cell line." Placenta (2013). Other ;Rabbit. 24112823
- **[IF=1.69]** Wang et al. Angiotensin 1-7 ameliorates caerulein-induced inflammation in pancreatic acinar cells by downregulating Toll-like receptor 4/nuclear factor-κB expression. (2018) Mol.Med.Rep. 17:3511-3518 ICC ;Rabbit. 29286117