bs-20786R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

WNT4 Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GenelD: 54361 **SWISS:** P56705

Target: WNT4

Immunogen: KLH conjugated synthetic peptide derived from human WNT4:

101-200/351.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Ligand for members of the frizzled family of seven transmembrane

receptors. Probable developmental protein. May be a signaling molecule which affects the development of discrete regions of tissues. Is likely to signal over only few cell diameters (By similarity). Overexpression may be associated with abnormal

proliferation in human breast tissue.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human, Mouse


(predicted: Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW.: 37 kDa


Subcellular Secreted ,Extracellular

Location: matrix

VALIDATION IMAGES -



Sample: Lane 1: MCF-7 (Human) Cell Lysate at 30 ug Lane 2: MDA-MB-231 (Human) Cell Lysate at 30 ug Lane 3: A431 (Human) Cell Lysate at 30 ug Lane 4: U2os (Human) Cell Lysate at 30 ug Primary: Anti-WNT4 (bs-20786R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 37 kD Observed band size: 47 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (WNT4) Polyclonal Antibody, Unconjugated (bs-20786R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

• [IF=2.626] Zhao, Gang. et al. LINC02381, a sponge of miR-21, weakens osteogenic differentiation of hUC-MSCs through KLF12-mediated Wnt4 transcriptional repression. 2021 Nov 15 WB; Mouse, Human. 34778905