

**bsm-60639M****[ Primary Antibody ]****GART Mouse mAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

|   |                             |   |
|---|-----------------------------|---|
| <b>Host:</b> Mouse  | <b>Isotype:</b> IgG1, Kappa | <b>Applications:</b> WB (1:500-1000)<br><b>ICC/IF</b> (1:100-500)<br><br><b>Reactivity:</b> Human (predicted: Rat)<br><br><br><b>Predicted MW.:</b> 107 kDa<br><br><b>Subcellular Location:</b> Cytoplasm |
| <b>Clonality:</b> Monoclonal  | <b>CloneNo.:</b> 1D14       |   |
| <b>GeneID:</b> 2618   | <b>SWISS:</b> P22102        |   |
| <b>Target:</b> GART   |                             |   |
| <b>Purification:</b> affinity purified by Protein A   |                             |   |
| <b>Concentration:</b> 1mg/ml  |                             |   |
| <b>Storage:</b> PBS, Glycerol, BSA.<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.  |                             |   |
| <b>Background:</b> Purines are critical for energy metabolism, cell signaling and cell reproduction and also function as precursors for coenzymes, energy transfer molecules, regulatory factors and proteins involved in RNA and DNA synthesis. GART (GAR transformylase), also referred to as AIRS, GARS, PAIS, PGFT, PRGS or GARTF, is 1,010 amino acids in length and is a key folate-dependent trifunctional enzyme with phosphoribosylglycinamide formyltransferase, phosphoribosylglycinamide synthetase and AICAR (phosphoribosylaminoimidazole synthetase) activity required for de novo purine biosynthesis. Cancer cells require considerable amounts of purines to sustain their accelerated growth and GART is, therefore, a target for cancer chemotherapy. GART is highly conserved in vertebrates. Two isoforms of GART are expressed due to alternative splicing events. |                             |   |

**— VALIDATION IMAGES —**

Sample: Lane 1: Human HeLa cell lysates Lane 2: Human A431 cell lysates Lane 3: Human HepG2 cell lysates Primary: Anti-GART (bsm-60639M) at 1/1000 dilution Secondary: IRDye800CW Goat Anti- Mouse IgG at 1/20000 dilution Predicted band size: 107 kDa Observed band size: 55,130 kDa