## bs-12678R

## [ Primary Antibody ]


## TMX3 Rabbit pAb




www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

	400-901-9800
<b>lsotype:</b> IgG	Applications: WB (1:500-2000)
	IHC-P (1:100-500) IHC-F (1:100-500)
SWISS: Q96JJ7	<b>IF</b> (1:100-500)
	Reactivity: Human, Mouse
tic peptide derived from human TMX3: smic >	
ein A	
	Predicted MW.: <sup>49 kDa</sup>
1% BSA, 0.02% Proclin300 and 50% : -20°C for one year. Avoid repeated	Subcellular Location: Cytoplasm
	SWISS: Q96JJ7 tic peptide derived from human TMX3: mic > ein A 1% BSA, 0.02% Proclin300 and 50%

## – VALIDATION IMAGES



Sample: Lung (Mouse) Lysate at 40 ug U251 (Human) Lysate at 40 ug Primary: Anti-TMX3 (bs-12678R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 49 kD Observed band size: 49 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TMX3) Polyclonal Antibody, Unconjugated (bs-12678R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.