bs-12679R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

VAC14 Rabbit pAb

GenelD: 55697 SWISS: Q08AM6

Target: VAC14

Immunogen: KLH conjugated synthetic peptide derived from human VAC14:

1-100/782.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The content of phosphatidylinositol 3,5-bisphosphate

(PtdIns(3,5)P2) in endosomal membranes changes dynamically with fission and fusion events that generate or absorb intracellular transport vesicles. VAC14 is a component of a trimolecular complex that tightly regulates the level of PtdIns(3,5)P2. Other components of this complex are the PtdIns(3,5)P2-synthesizing enzyme PIKFYVE (MIM 609414) and the PtdIns(3,5)P2 phosphatase FIG4 (MIM 609390). VAC14 functions as an activator of PIKFYVE (Sbrissa et al.,

2007 [PubMed 17556371]).[supplied by OMIM, Feb 2010]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Mouse (predicted: Human,


Rat, Rabbit, Pig, Cow, Zebrafish, Chicken, Dog,

Horse)

Predicted MW.: 88 kDa

Subcellular Cytoplasm

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (VAC14) Polyclonal Antibody, Unconjugated (bs-12679R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.