## bs-5732R

## [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

# **NEK2 Rabbit pAb**

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 4751 **SWISS:** P51955

Target: NEK2

Immunogen: KLH conjugated synthetic peptide derived from human NEK2:

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Protein kinase that is involved in mitotic regulation. Integral component of the mitotic spindle-assembly checkpoint which is necessary for proper chromosome segregation during metaphaseanaphase transition. Required for association of MAD2L1 to kinetochore. Phosphorylates SGOL1. May have a role at the G2-M transition. May also play a role in meiosis. Isoform 1 but not isoform 2 appears to play a role in centrosome splitting. Isoform 1 phosphorylates and activates NEK11 in G1/S-arrested cells. Isoform 2, which is not present in the nucleolus, does not.

Applications: WB (1:500-2000)

400-901-9800

Reactivity: Human (predicted: Mouse,


Rat, Rabbit, Pig, Cow, Dog,

Horse)

Predicted 52 kDa MW.:

Subcellular Cytoplasm , Nucleus

## VALIDATION IMAGES -



Sample:Lovo Cell (Human) Lysate at 40 ug Primary: Anti-NEK2(bs-5732R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-RabbitIgG at 1/20000 dilution Predicted band size: 52kD Observed band size: 52kD

### — SELECTED CITATIONS -

- [IF=8.378] Roberts MS et al. LIN9 and NEK2 are core regulators of mitotic fidelity that can be therapeutically targeted to overcome taxane resistance. Cancer Res. 2020 Feb 13. WB; Human. 32054769
- [IF=2.52] Zhong, Xinwen, et al. "Examining Nek2 as a better proliferation marker in non-small cell lung cancer prognosis." Tumor Biology (2014): 1-8. IHC ;="Human". 24763826
- [IF=0.99] Zhong, Xinwen, et al. "Aberrant expression of NEK2 and its clinical significance in non-small cell lung cancer." Oncology Letters. IHC;="Human". 25202351
- [IF=1.41] Zeng, Yan-Ru, et al. "Overexpression of NIMA-related kinase 2 is associated with progression and poor prognosis of prostate cancer." BMC Urology 15.1 (2015): 90. WB ;="Human". 26320076