

bs-10949R**[Primary Antibody]****ER81 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit Clonality: Polyclonal GeneID: 2115 Target: ER81 Immunogen: KLH conjugated synthetic peptide derived from human ER81: 51-150/477. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Several members of the Ets gene family encode sequence-specific DNA binding proteins that recognize DNA sequences with a centrally located 5'-GGAA-3' element. All of the Ets proteins recognize the same central core sequence but each protein interacts with unique sequences that flank this core. PEA3 binds the motif 5'-AGGAAG-3', while ER81 (also designated ETV1) binds the motif 5'-CGGAA/T-3'. PEA3 is expressed at readily detectable levels in cells of epithelial and fibroblastic origin. Unlike other members of the Ets family, including Ets-1 and Ets-2, PEA3 is not expressed in hematopoietic cells. ER81 is highly expressed in brain, testis, lung and heart. ER81 is also moderately expressed in spleen, pancreas, colon and small intestine. During development, ER81, PEA3 and ERM display unique expression patterns which suggest these transcriptional factors play an important role in organogenesis. ERK-1 activates ER81 transcriptional activity, while MAPKAP kinase 2 inhibits ER81.	Isotype: IgG SWISS: P50549	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
		Reactivity: Mouse, Rat (predicted: Human, Rabbit, Pig, Sheep, Cow, Dog, Horse)
		Predicted MW.: 55 kDa
		Subcellular Location: Nucleus

— VALIDATION IMAGES —

Sample: Lane 1: Mouse Cerebellum tissue lysates
Lane 2: Mouse Heart tissue lysates Primary: Anti-ER81 (bs-10949R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kDa Observed band size: 53 kDa



Paraformaldehyde-fixed, paraffin embedded (mouse cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ER81) Polyclonal Antibody, Unconjugated (bs-10949R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ER81) Polyclonal Antibody, Unconjugated (bs-10949R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=4.6]** Tianci Han. et al.SUMO modified ETV1 promotes M2-polarized tumor-associated macrophage infiltration and cancer progression by facilitating CCL2 transcription in esophageal squamous cell carcinoma cells.CANCER IMMUNOLOGY

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

