bs-0294D-BF555

## [ Secondary Antibodies ]

## Donkey Anti-Goat IgG H&L, BF555 conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| – DATASHEET ––––––  |  | 400-901-9800  |
|---|--|---|
| Host: Donkey  | Isotype: IgG   | Applications: IF (1:100-1000)<br>Flow-Cyt (1:100-1000)  |
| Clonality: Polyclonal<br>Target: Donkey Anti-Goat IgG<br>Purification: affinity purified by Pro<br>Concentration: 2.0 mg/ml<br>Storage: 10 mM TBS (pH=7.4) v<br>glycerol.   |  | ICC/IF (1:100-1000)<br>Excitation Spectrum: 555nm<br>Emission spectrum: 572nm<br>Reactivity: Goat |
| Store at -20°C for one<br>Background: Immunoglobulin G (Ig<br>serum with normal le<br>is important for our d<br>molecules are produc<br>adaptive immune res<br>functions; to bind to t<br>recruit other cells and<br>variability of the IgG p | vels between 8-17 mg/mL in adult bl<br>efence against microorganisms and<br>ed by B lymphocytes as a part of our<br>ponse. The IgG molecule has two sep<br>he pathogen that elicited the respor<br>I molecules to destroy the antigen. T<br>pool is generated by somatic recomb<br>ecificities in an individual at a given | teins in<br>lood. IgG<br>the<br>r<br>parate<br>nse and to<br>The<br>pination                      |