bs-20658R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

FSH receptor Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 2492 SWISS: P23945

Target: FSH receptor

Immunogen: KLH conjugated synthetic peptide derived from human FSH

receptor: 21-100/695. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene belongs to family 1 of G-protein

coupled receptors. It is the receptor for follicle stimulating hormone and functions in gonad development. Mutations in this

gene cause ovarian dysgenesis type 1, and also ovarian hyperstimulation syndrome. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Mar 2010]

Applications: WB (1:500-2000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) Flow-Cyt (1ug/Test)


Reactivity: Human, Mouse, Rat

(predicted: Pig, Sheep, Cow, Dog, Horse)


Predicted 78 kDa MW.:

Subcellular Location: Cell membrane

VALIDATION IMAGES




Sample: Ovary (mouse) Lysate at 40 ug Primary: Anti- FSH receptor (bs-20658R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 78kD Observed band size: 78kD



Paraformaldehyde-fixed, paraffin embedded (rat ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FSH receptor) Polyclonal Antibody, Unconjugated (bs-20658R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CBFb) Polyclonal Antibody, Unconjugated (bs-20658R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control (blue line): MCF7 (fixed with 70% methanol overnight at 4°C). Primary Antibody (green line): Rabbit Anti-FSH receptor antibody (bs-20658R), Dilution: 0.2µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat antirabbit IgG-PE, Dilution: 1µg/test.

- [IF=6.208] Sijie Fan. et al. Seasonal Change in Adiponectin Associated with Ovarian Morphology and Function in Wild Ground Squirrels (Citellus dauricus Brandt). INT J MOL SCI. 2022 Jan;23(23):14698 IHC; Squirrel. 36499026
- [IF=6] Yu Cai. et al. β-Nicotinamide Mononucleotide Reduces Oxidative Stress and Improves Steroidogenesis in Granulosa Cells Associated with Sheep Prolificacy via Activating AMPK Pathway. ANTIOXIDANTS-BASEL. 2025 Jan;14(1):34 IF; Sheep. 10.3390/antiox14010034
- [IF=5.108] Yang Z et al.

 A pilot study on polycystic ovarian syndrome caused by neonatal exposure to tributyltin and bisphenol A in rats.

 Chemosphere. 2019 Sep;231:151-160. WB;Rat. 31129395
- [IF=4.627] Wang Q et al.

 Heat shock pretreatment improves mesenchymal stem cell viability by heat shock proteins and autophagy to prevent cispl atin-induced granulosa cell apoptosis. Stem Cell Res Ther. 2019 Nov 26;10(1):348. ICC; Rat. 31771642
- [IF=4.982] Luo, Qianqian. et al. hUCMSCs reduce theca interstitial cells apoptosis and restore ovarian function in premature ovarian insufficiency rats through regulating NR4A1-mediated mitochondrial mechanisms. REPROD BIOL ENDOCRIN. 2022 Dec;20(1):1-14 IF; Rat. 35986315