

bs-9449R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

TET2 Rabbit pAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200) Flow-Cyt (2ug/Test) ICC/IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Pig, Sheep, Cow, Chicken, Dog, GuineaPig, Horse) Predicted MW.: 224 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 54790	SWISS: Q6N021	
Target: TET2		
Immunogen: KLH conjugated synthetic peptide derived from human TET2: 1101-1300/2002.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: TET2 is a 2,002 amino acid protein that is expressed in a variety of tissues, including brain, kidney, heart, lung, muscle and stomach, and exists as three alternatively spliced isoforms. Murine TET2 is also known as protein Ayu17-449 and is thought to play a role in proper kidney development and overall kidney function, as well as in hormone secretion throughout the body. The gene encoding human TET2 maps to chromosome 4 and the gene encoding mouse TET2 maps to chromosome 3. Chromosome 4 encodes nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes. Defects in some of the genes located on chromosome 4 are associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease. Murine chromosome 3 houses over 1,300 genes, some of which express alcohol dehydrogenases (ADHs), sodium channel modifiers (SCNMs), interleukins (ILs) and Insulin receptor-related (IRR) proteins. Defects in chromosome 3-localized genes are associated with hereditary congenital facial paresis (HCFP), increased susceptibility to spontaneous colitis, HIV-1-associated nephropathy, decreased renal vascular health and malignant sporadic pancreatic endocrine tumors.		

— VALIDATION IMAGES —


Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TET2) Polyclonal Antibody, Unconjugated (bs-9449R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TET2) Polyclonal Antibody, Unconjugated (bs-9449R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



MCF7 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (TET2) polyclonal Antibody, Unconjugated (bs-9449R) 1:25, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (black line) :MCF-7. Primary Antibody (green line): Rabbit Anti-TET2 antibody (bs-9449R) Dilution:2ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=6.61]** Bras et al. Adventitial Sca1+ Cells Transduced With ETV2 Are Committed to the Endothelial Fate and Improve Vascular Remodeling After Injury. (2018) Arterioscler.Thromb.Vasc.Biol. 38:232-244 Other ;. 29191922
- **[IF=2.9]** Fuyi Luo. et al. Simultaneous blastic plasmacytoid dendritic cell neoplasm and myelofibrosis: A case report. ONCOL LETT. 2024 May;27(5):1-14 IHC ;Human. 38586204