bs-5210R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-TEM8 (Tyr382) Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 84168 SWISS: Q9H6X2

Target: TEM8 (Tyr382)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

ANTXR1 around the phosphorylation site of Tyr382: AS(p-Y)YG.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a type I transmembrane protein and is a tumorspecific endothelial marker that has been implicated in colorectal cancer. The encoded protein has been shown to also be a docking protein or receptor for Bacillus anthracis toxin, the causative agent of the disease, anthrax. The binding of the protective antigen (PA) component, of the tripartite anthrax toxin, to this receptor protein mediates delivery of toxin components to the cytosol of cells. Once inside the cell, the other two components of anthrax toxin, edema factor (EF) and lethal factor (LF) disrupt normal cellular processes. Three alternatively spliced variants that encode different protein isoforms have been described. [provided by RefSeq, Oct 2008]

Applications: WB (1:500-2000)

Reactivity: Mouse (predicted: Human,


Rat, Rabbit, Pig, Sheep,

Chicken)


Predicted 45 kDa MW.:

Subcellular Location: Cell membrane

VALIDATION IMAGES



Sample: Bladder (Mouse) Lysate at 40 ug Primary: Anti-phospho-TEM8 (Tyr382)(bs-5210R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 45 kD Observed band size: 45 kD



Sample: Uterus (Mouse) Lysate at 40 ug Primary: Anti-phospho-TEM8 (Tyr382)(bs-5210R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 45 kD Observed band size: 45 kD