

bs-17341R**[Primary Antibody]****hnRNP A3 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

| | | |
|---|---|---|
| Host: Rabbit Clonality: Polyclonal GeneID: 220988 Target: hnRNP A3 Immunogen: KLH conjugated synthetic peptide derived from human hnRNP A3: 61-160/378. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Heterogeneous nuclear ribonucleoproteins (hnRNPs) constitute a set of poly-peptides that contribute to mRNA transcription and pre-mRNA processing as well as mature mRNA transport to the cytoplasm and translation. They also bind heterogeneous nuclear RNA (hnRNA), which are the transcripts produced by RNA polymerase II. There are approximately 20 known hnRNP proteins, and their complexes are the major constituents of the spliceosome. The majority of hnRNP protein components are localized to the nucleus; however some shuttle between the nucleus and the cytoplasm. The A/B subfamily of hnRNPs include A1, A2/B1, A3 and A0, and in Xenopus, hnRNP A1, A2 and A3 are ubiquitously expressed throughout development as well as in adult tissues. hnRNP A1 and A2/B1 regulate the processing of pre-mRNA by directly antagonizing the association of various splicing factors and by influencing the splice site selection on pre-mRNA. The hnRNP A0 gene is distinct from the other A/B family members, and it encodes a low-abundance protein, which is implicated in mRNA stability. | Isotype: IgG SWISS: P51991 | Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) |
| | | Reactivity: Human, Mouse (predicted: Rat, Rabbit, Cow, Dog, Horse) |
| | | Predicted MW.: 40 kDa |
| | | Subcellular Location: Nucleus |

— VALIDATION IMAGES —

Tissue/cell: mouse intestine tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-hnRNP A3 Polyclonal Antibody, Unconjugated(bs-17341R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: HepG2 cells; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-hnRNP A3 Polyclonal Antibody, Unconjugated(bs-17341R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining