bsm-30041M

[Primary Antibody]

BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

human CD45RA Mouse mAb

- DATASHEET -

Host: Mouse IgG1, k

Clonality: Monoclonal CloneNo.: 45F1

GenelD: 5788 SWISS: P08575

Target: human CD45RA

Purification: affinity purified by Protein G

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4).

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a member of the protein

tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus belongs to receptor type PTP. This gene is specifically expressed in hematopoietic cells. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus


functions as a regulator of cytokine receptor signaling

Applications: Flow-Cyt (lug/Test)

Reactivity: Human

Subcellular Location: Cell membrane

VALIDATION IMAGES



Blank control:Jurkat. Primary Antibody (green line): Mouse Anti-human CD45RA antibody (bsm-30041M) Dilution: 1ug/Test; Secondary Antibody (white blue line): Goat anti-mouse IgG-FITC Dilution: 0.5ug/Test. Isotype control (orange line): Normal Rabbit IgG Protocol The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.