

bs-0449R**[Primary Antibody]****TSLP Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

DATASHEET


Host: Rabbit Clonality: Polyclonal GeneID: 85480 Target: TSLP Immunogen: KLH conjugated synthetic peptide derived from human TSP-1: 101-159/159. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Thymic stromal lymphopoietin (TSLP) is a novel cytokine that was found to promote the development of murine B cells in vitro. This protein, which is expressed in a number of tissues including heart, liver and prostate, prevented apoptosis and stimulated growth of the human acute myeloid leukemia (AML)-derived cell line MUTZ-3. Anti-interleukin (IL)-7 receptor antibodies (Abs) neutralized this effect indicating that TSLP binds to at least part of the IL-7 receptor complex. TSLP induced phosphorylation of signal transducer and activator of transcription (STAT)-5.	Isotype: IgG SWISS: Q969D9	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human Predicted MW.: 15 kDa Subcellular Location: Secreted
--	---	---

VALIDATION IMAGES

Sample: Lane1:K562 Cell Lysate at 30ug Lane2: Raji Cell Lysate at 30ug Primary: Anti-TSLP (bs-0449R) at 1:300 dilution; Secondary: HRP conjugated Goat-Anti-rabbit IgG(bs-0295G-HRP) at 1: 5000 dilution; Predicted band size: 15 kD
Observed band size: 17 kD



Tissue/cell: human liver carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-TSLP Polyclonal Antibody, Unconjugated(bs-0449R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TSLP) Polyclonal Antibody, Unconjugated (bs-0449R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

SELECTED CITATIONS

- **[IF=6.8]** Jana Zeitvogel. et al. Short-form thymic stromal lymphopoietin (sfTSLP) restricts herpes simplex virus infection of human primary keratinocytes. J MED VIROL. 2024 Sep;96(9):e29865 WB ;Human. 39233492
- **[IF=5.34]** Dian-Dong Hou. et al. Therapeutic effects of myricetin on atopic dermatitis in vivo and in vitro. PHYTOMEDICINE. 2022 Jul;102:154200 IHC ;Mouse. 35671605
- **[IF=3.483]** Shen Z et al. Insights into Roseburia intestinalis which alleviates experimental colitis pathology by inducing anti-inflammatory responses. J Gastroenterol Hepatol. 2018 Oct;33(10):1751-1760. IHC ;Mouse. 29532517

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- **[IF=2.493]** Min-jie Gong, et al. Melatonin reduces IL-33 and TSLP expression in human nasal epithelial cells by scavenging ROS directly. IMMUN INFLAMM DIS. 2023 Feb;11(2):e788 WB ;Human. 36840497
- **[IF=0.67]** Sakitani, Eri, et al. "Increased Expression of Thymic Stromal Lymphopoietin and Its Receptor in Kimuras Disease." ORL 77.1 (2015): 44-54. Other ;="". 25676453