

**bs-21215R**

**[ Primary Antibody ]**

## SLC23A2 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)  <b>Reactivity:</b> (predicted: Human, Mouse, Rat, Rabbit)  <b>Predicted MW.:</b> 70 kDa  <b>Subcellular Location:</b> Cell membrane
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 9962	<b>SWISS:</b> Q9UGH3	
<b>Target:</b> SLC23A2		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human SLC23A2: 551-650/650.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The absorption of vitamin C into the body and its distribution to organs requires two sodium-dependent vitamin C transporters. This gene encodes one of the two required transporters and the encoded protein accounts for tissue-specific uptake of vitamin C. Previously, this gene had an official symbol of SLC23A1. [provided by RefSeq, Jul 2008]		