bs-8770R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Beta Actin (CAEEL) Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

Target: Beta Actin (CAEEL)

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.


Applications: WB (1:500-2000)

Reactivity: Caenorhabditis Elegans

Predicted MW.: 42 kDa

Subcellular Location: Cytoplasm

VALIDATION IMAGES



Sample: Lane 1: Caenorhabditis Elegans tissue Lysate Primary: Anti-Beta Actin (CAEEL) (bs-8770R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 42 kD Observed band size: 42 kD

— SELECTED CITATIONS ——

- [IF=4.162] Zhen-Ni Liu. et al. Blockade of the P2Y2 Receptor Attenuates Alcoholic Liver Inflammation by Targeting the EGFR-ERK1/2 Signaling Pathway. DRUG DES DEV THER. Drug Des Dev Ther. 2022 Apr;16:1107-1120 WB; Mouse.
- [IF=4.2] Muxin Yu. et al. Neutrophil extracellular traps-induced pyroptosis of liver sinusoidal endothelial cells exacerbates intrahepatic coagulation in cholestatic mice. BBA-MOL BASIS DIS. 2025 Mar;1871:167700 WB; Mouse. 39914029
- [IF=2.794] Ji, Shanghong. et al. PKA/ATGL signaling pathway is involved in ER stress-mediated lipolysis in adipocytes of grass carp (Ctenopharyngodon idella). FISH PHYSIOL BIOCHEM. 2022 Apr;:1-9 WB; Fish. 35460470
- [IF=2.74] Rui Feng. et al. Progesterone regulates inflammation and receptivity of cells via the NF-kB and LIF/STAT3 pathways. THERIOGENOLOGY. Theriogenology. 2022 Jul;186:50 WB;Bovine. 35430548
- [IF=2.72] Liu Yang. et al. Protective effect of isopsoralen on UVB-induced injury in HaCaT cells via the ER and p38MAPK signaling pathways. J FOOD BIOCHEM. 2022 Apr 12 WB; Human. 35415935