bs-0294M-PE

## [ Secondary Antibodies ]

## Mouse Anti-Goat IgG H&L, PE conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET ———		400-901-9800
Host: Mouse	<b>lsotype:</b> IgG	Applications: IF (1:100-1000) Flow-Cyt (1:100-1000)
Clonality: Polyclonal		ICC/IF (1:100-1000)
Target: Mouse Anti-Goat IgG H&L Purification: affinity purified by Protein G Concentration: 2.0 mg/ml		Excitation Spectrum: 496/564nm Emission spectrum: 578nm
		Reactivity: Goat
glycerol.	with 1% BSA, 0.03% Proclin300 and 5 9 year. Avoid repeated freeze/thaw cy	
serum with normal le is important for our o molecules are produ adaptive immune res functions; to bind to recruit other cells an variability of the IgG	gG), is one of the most abundant protected between 8-17 mg/mL in adult blefence against microorganisms and iced by B lymphocytes as a part of our ponse. The IgG molecule has two septhe pathogen that elicited the respond molecules to destroy the antigen. To pool is generated by somatic recomb becificities in an individual at a given to be 1011 variants.	ood. IgG the r parate use and to he ination