bs-0293R-PE-CY5

## [ Secondary Antibodies ]

## Rabbit Anti-Rat IgG H&L, PE-CY5 conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit Clonality: Polyclonal	<b>lsotype:</b> lgG	Applications: IF (1:100-1000) Flow-Cyt (1:100-1000) ICC/IF (1:100-1000)
glycerol. Store at -20°C for or <b>Background:</b> Immunoglobulin G serum with normal is important for our molecules are produ adaptive immune re functions; to bind to recruit other cells ar variability of the IgG	Protein A with 1% BSA, 0.03% Proclin300 and 5 we year. Avoid repeated freeze/thaw cy (IgG), is one of the most abundant pro- levels between 8-17 mg/mL in adult bl defence against microorganisms and uced by B lymphocytes as a part of our esponse. The IgG molecule has two sep the pathogen that elicited the respor- nd molecules to destroy the antigen. T pool is generated by somatic recomb specificities in an individual at a given	Excitation Spectrum: 488nm Emission spectrum: 670nm Reactivity: Rat weles. teins in lood. IgG the r parate nse and to The pination