bs-0293R-PE-Cy7

## [ Secondary Antibodies ]

## Rabbit Anti-Rat IgG H&L, PE-Cy7 conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit Clonality: Polyclonal	<b>lsotype:</b> IgG	Applications: IF (1:100-1000) Flow-Cyt (1:100-1000) ICC/IF (1:100-1000)
glycerol. Store at -20°C for on <b>Background:</b> Immunoglobulin G ( serum with normal l is important for our molecules are produ adaptive immune re functions; to bind to recruit other cells ar variability of the IgG	rotein A with 1% BSA, 0.03% Proclin300 and 5 e year. Avoid repeated freeze/thaw cy lgG), is one of the most abundant pro evels between 8-17 mg/mL in adult bl defence against microorganisms and uced by B lymphocytes as a part of ou sponse. The IgG molecule has two sep the pathogen that elicited the respon id molecules to destroy the antigen. T pool is generated by somatic recomb pecificities in an individual at a given	Excitation Spectrum: 488nm Emission spectrum: 780nm Reactivity: Rat 50% ycles. oteins in lood. IgG the parate nse and to The bination