

bs-9163R**[Primary Antibody]****RNF17 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Rat (predicted: Human, Mouse, Sheep, Cow, Dog, Horse) Predicted MW.: 185 kDa Subcellular Location: Cytoplasm ,Nucleus
Clonality: Polyclonal		
GeneID: 56163	SWISS: Q9BXT8	
Target: RNF17		
Immunogen: KLH conjugated synthetic peptide derived from human RNF17: 1031-1200/1623.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. RNF17 (ring finger protein 17) or tudor domain-containing protein 4, TDRD4, SPATA23, Mmip-2 or FLJ11045, is a testis-specific protein and a novel key regulator of spermiogenesis containing 1,623 amino acids. By distributing Mad proteins to the cytoplasm, RNF17 regulates the transcriptional activity of c-Myc. Although showing localization in the nucleus, RNF17 is predominantly observed in cytoplasm and is a component of a novel nuage found in male germ cells. The gene encoding RNF17 maps to human chromosome 13q12.12 and encodes one RING-type zinc finger and four tudor domains. As a result of alternative splice events, five RNF17 isoforms exist.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF17) Polyclonal Antibody, Unconjugated (bs-9163R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining