

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **HOXB7** Rabbit pAb

Catalog Number: bs-17364R

Target Protein: HOXB7
Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:100-500),

ELISA (1:5000-10000)

Reactivity: (predicted:Human, Mouse, Rat, Sheep, Chicken, Horse)

Predicted MW: 24 kDa
Subcellular Nucleus

Locations:

Entrez Gene: 3217 Swiss Prot: P09629

Source: KLH conjugated synthetic peptide derived from human HOXB7: 101-200/217.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The Hox proteins play a role in development and cellular differentiation by regulating

downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. The mammalian HOX gene complex consists of 39 genes that are located on four linkage groups, which are dispersed over four chromosomes. A segment of the HoxB7 proximal promoter drives renal expression of reporter genes specifically in the ureteric bud and collecting ducts. Expression levels of HoxB7 are lower in lymph node metastasis-positive cancer tissues than negative cancer tissues. These results suggest that aberrant expression of HOX genes is related to the development of breast cancer and

malignant behavior of cancer cells.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=3.5] Yuanhui Wang. et al.A pan-cancer analysis of homeobox family: expression characteristics and latent significance in prognosis

and immune microenvironment.front oncol.2025 Feb 6:15:1521652. IHC; Human . 39980564	