

**bs-5226R****[ Primary Antibody ]****phospho-BRIP1/BACH1 (Ser990) Rabbit pAb****Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


## — DATASHEET —

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 83990</p> <p><b>Target:</b> phospho-BRIP1/BACH1 (Ser990)</p> <p><b>Immunogen:</b> KLH conjugated Synthesised phosphopeptide derived from human BRIP1 around the phosphorylation site of Ser990: ST(p-S)PT.</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> This gene encodes a transcription factor that belongs to the cap'n'collar type of basic region leucine zipper factor family (CNC-bZip). The encoded protein contains broad complex, tramtrack, bric-a-brac/poxvirus and zinc finger (BTB/POZ) domains, which is atypical of CNC-bZip family members. These BTB/POZ domains facilitate protein-protein interactions and formation of homo- and/or hetero-oligomers. When this encoded protein forms a heterodimer with MafK, it functions as a repressor of Maf recognition element (MARE) and transcription is repressed. Multiple alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, May 2009].</p>	<p><b>Isotype:</b> IgG</p> <p><b>SWISS:</b> Q9BX63</p> <p><b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p><b>Reactivity:</b> Human, Mouse (predicted: Sheep, Cow)</p> <p><b>Predicted MW.:</b> 141 kDa</p> <p><b>Subcellular Location:</b> Nucleus</p>
--	---

## — VALIDATION IMAGES —



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-BRIP1/BACH1 (Ser990)) Polyclonal Antibody, Unconjugated (bs-5226R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: Mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-BACH 1 Polyclonal Antibody, Unconjugated(bs-5226R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining