

**bs-20646R****[ Primary Antibody ]****RANKL/CD254 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**DATASHEET**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>Flow-Cyt</b> (1ug/Test)
<b>Clonality:</b> Polyclonal		<b>Reactivity:</b> Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Zebrafish, Dog, Horse)
<b>GeneID:</b> 8600	<b>SWISS:</b> Q14788	<b>Predicted MW.:</b> 35 kDa
<b>Target:</b> RANKL/CD254		<b>Subcellular Location:</b> Secreted ,Cell membrane Cytoplasm
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human RANKL/CD254: 41-140/317.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> This gene encodes a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. This protein was shown to be a dendritic cell survival factor and is involved in the regulation of T cell-dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor-associated factor (TRAF) 6, which indicated this protein may have a role in the regulation of cell apoptosis. Targeted disruption of the related gene in mice led to severe osteopetrosis and a lack of osteoclasts. The deficient mice exhibited defects in early differentiation of T and B lymphocytes, and failed to form lobulo-alveolar mammary structures during pregnancy. Two alternatively spliced transcript variants have been found. [provided by RefSeq, Jul 2008].		

**VALIDATION IMAGES**

Sample: Lung (Mouse) Lysate at 40 ug Primary:  
Anti-CD254 (bs-20646R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at  
1/20000 dilution Predicted band size: 35 kD  
Observed band size: 37/50 kD



Sample: Lymph node (Mouse) Lysate at 40 ug  
Thymus (Mouse) Lysate at 40 ug Lymph node  
(Rat) Lysate at 40 ug Primary: Anti-RANKL'CD254  
(bs-20646R) at 1/500 dilution Secondary:  
IRDye800CW Goat Anti-Rabbit IgG at 1/20000  
dilution Predicted band size: 37'50 kD Observed  
band size: 50 kD



Blank control:HL-60. Primary Antibody (green  
line): Rabbit Anti-RANKL/CD254 antibody  
(bs-20646R) Dilution: 1μg /10<sup>6</sup> cells; Isotype  
Control Antibody (orange line): Rabbit IgG .  
Secondary Antibody : Goat anti-rabbit IgG-AF488  
Dilution: 1μg /test. Protocol The cells were fixed  
with 4% PFA (10min at room temperature)and  
then permeabilized with 0.1% PBST for 20 min at  
room temperature. The cells were then  
incubated in 5%BSA to block non-specific  
protein-protein interactions for 30 min at room  
temperature .Cells stained with Primary  
Antibody for 30 min at room temperature. The  
secondary antibody used for 40 min at room  
temperature. Acquisition of 20,000 events was  
performed.

## — SELECTED CITATIONS —

---

- **[IF=0]** Francesca Punzo et al. Can Denosumab be used in combination with Doxorubicin in Osteosarcoma?. Oncotarget . 2020 Jul 14;11(28):2763-2773. WB ;Human. 32733647