

bsm-30235M**[Primary Antibody]****Human Bcl-2 Mouse mAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Mouse	Isotype: Mouse IgG2b,k	Applications: Flow-Cyt (1ug/Test) ICC/IF (1:50-200)
Clonality: Monoclonal	CloneNo.: 8B4	
Target: Human Bcl-2		Reactivity: Human
Purification: affinity purified by Protein G		
Storage: 0.01M TBS (pH7.4). Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 26 kDa
Background: The Bcl-2 gene was isolated at the chromosomal breakpoint of t(14;18)-bearing follicular B cell lymphomas(1,2).Bcl-2 blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal (3,5).Bcl-2 appears to function in several subcellular locations yet lacks any known motifs that would confer insight into its mechanism of action (6,7).A more recently identified protein,designated Bax p21(i.e., Bcl-associated X protein),has extensive amino acid homology with Bcl-2 and both homodimerizes and forms heterodimers with Bcl-2(8). Overexpression of Bax accelerates apoptotic death induced by cytokine deprivation in an IL-3 dependent cell line and Bax also counters the death repressor activity of Bcl-2(8).		Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus

— VALIDATION IMAGES —

4% Paraformaldehyde-fixed HeLa (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (Bcl-2) monoclonal Antibody, unconjugated (bsm-30235M) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Mouse IgG antibody (green, bs-60296G-BF488) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.



The HeLa (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.).Primary Antibody (green):Mouse Anti-Bcl-2 antibody (bsm-30235M): 1 µg/10⁶ cells; Secondary Antibody (white blue): Goat anti-Mouse IgG-BF488(bs-60296G-BF488): 1 µg/test. Blank control (black): PBS. Acquisition of 20,000 events was performed.