

**bs-16408R**

**[ Primary Antibody ]**

## ZRSR2 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 8233</p> <p><b>Target:</b> ZRSR2</p> <p><b>Immunogen:</b> KLH conjugated synthetic peptide derived from human ZRSR2: 351-450/482.</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> This gene encodes an essential splicing factor. The encoded protein associates with the U2 auxiliary factor heterodimer, which is required for the recognition of a functional 3' splice site in pre-mRNA splicing, and may play a role in network interactions during spliceosome assembly. [provided by RefSeq, Jul 2008]</p>	<p><b>Isotype:</b> IgG</p> <p><b>SWISS:</b> Q15696</p>	<p><b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ICC/IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)</p> <p><b>Reactivity:</b> (predicted: Human)</p> <p><b>Predicted MW.:</b> 58 kDa</p> <p><b>Subcellular Location:</b> Nucleus</p>
--	--	---