

**bs-3402R****[ Primary Antibody ]****phospho-INPPL1 (Ser576) Rabbit pAb****Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn



techsupport@bioss.com.cn

400-901-9800

## — DATASHEET —

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 3636</p> <p><b>Target:</b> INPPL1 (Ser576)</p> <p><b>Immunogen:</b> KLH conjugated Synthesised phosphopeptide derived from human SHIP1 around the phosphorylation site of Ser576: QN(p-Y)LD.</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> The steady state of protein tyrosyl phosphorylation in cells is regulated by the opposing action of tyrosine kinases and protein tyrosine phosphatases (PTPs). Several groups have independently identified a non transmembrane PTP, designated SHPTP1 (also known as PTP1C, HCP and SHP), which is primarily expressed in hematopoietic cells and characterized by the presence of two SH2 domains N terminal to the PTP domain. A second and much more widely expressed PTP with SH2 domains, SHPTP2 (also designated PTP1D and Syp), has been identified. SHP2 is a protein tyrosine phosphatase that is widely expressed and plays a regulatory role in various cell signaling events that are important for many cell functions, such as mitogenic activation, metabolic control, transcription regulation, and cell migration.</p>	<p><b>Isotype:</b> IgG</p> <p><b>SWISS:</b> O15357</p>	<p><b>Applications:</b> WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p><b>Reactivity:</b> Mouse, Rat (predicted: Human, Rabbit, Pig, Cow, Dog, Horse)</p> <p><b>Predicted MW.:</b> 139 kDa</p> <p><b>Subcellular Location:</b> Cell membrane ,Cytoplasm</p>
--	--	--

## — VALIDATION IMAGES —



Sample: Uters (Mouse) Lysate at 30 ug Primary: Anti- p-INPPL1 (Ser576) (bs-3402R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/10000 dilution Predicted band size: 139 kD Observed band size: 135kD

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (INPPL1) Polyclonal Antibody, Unconjugated (bs-3402R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.