bs-2808R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Jak3 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 3718 SWISS: P52333

Target: Jak3

Immunogen: KLH conjugated synthetic peptide derived from human Jak3:

601-700/1124.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a member of the Janus kinase

(JAK) family of tyrosine kinases involved in cytokine receptormediated intracellular signal transduction. It is predominantly expressed in immune cells and transduces a signal in response to its activation via tyrosine phosphorylation by interleukin receptors. Mutations in this gene are associated with autosomal SCID (severe combined immunodeficiency disease). [provided by RefSeq, Jul

2008]

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1ug/test)

Reactivity: Human, Mouse, Rat

(predicted: Pig, Cow, Dog,


Horse)

Predicted MW.: 125 kDa

Subcellular

Location: Cytoplasm


- VALIDATION IMAGES -




Sample: SP2/0 Cell (Mouse) Lysate at 40 ug Jurkat Cell (Human) Lysate at 40 ug K562 Cell (Human) Lysate at 40 ug Primary: Anti- Jak3 (bs-2808R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 125 kD Observed band size: 125 kD




Sample: Lane 1: Mouse Thymus tissue lysates Lane 2: Rat Lymph node tissue lysates Lane 3: Human Jurkat cell lysates Lane 4: Human Hela cell lysates Lane 5: Human Molt-4 cell lysates Primary: Anti-Jak3 (bs-2808R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 125 kDa Observed band size: 115 kDa



Tissue/cell: human lung carcinoma: 4% Paraformaldehyde-fixed and paraffinembedded: Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Jak3 Polyclonal Antibody, Unconjugated(bs-2808R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control (black line) :Molt4. Primary Antibody (green line): Rabbit Anti-Jak3 antibody (bs-2808R) Dilution:1ug/Test; Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line): Normal Rabbit IgG Protocol The



Blank control:Molt-4. Primary Antibody (green line): Rabbit Anti-Jak3 antibody (bs-2808R) Dilution: 0.2µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-PE Dilution: 0.2µg /test. Protocol The cells were fixed with 4% PFA

cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

(10min at room temperature) and then permeabilized with 20% PBST for 20 min at-20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=17.4] Hui Yan. et al. Elaboration a ROS-responsive darutigenol prodrug nanoassemblies for inflammatory arthritis treatment. NANO TODAY. 2024 Apr;55:102220 WB ;Mouse. 10.1016/j.nantod.2024.102220
- [IF=14.1] Jiayu Wang. et al. Discovery of Natural Compound α-Hederin via Large-Scale Screening as a Targeted JAK/STAT3 Inhibitor for Ovarian Cancer Therapy. ADV SCI. 2025 Jul;:e17278 WB; Human. 40667652
- [IF=6.291] Changjiang Liu. et al. Cypermethrin triggers YY1-mediated testosterone biosynthesis suppression. Ecotox Environ Safe. 2021 Dec;225:112792 WB;Rat. 10.1016/j.ecoenv.2021.112792
- [IF=5.195] Yong Wang. et al. An integrated network pharmacology approach reveals that Darutigenol reduces inflammation and cartilage degradation in a mouse collagen-induced arthritis model by inhibiting the JAK-STAT3 pathway. J ETHNOPHARMACOL. 2023 May;:116574 WB; Mouse. 37160212
- [IF=5.4] Ting Xiao. et al. Ameliorative effect of Alangium chinense (Lour.) Harms on rheumatoid arthritis by reducing autophagy with targeting regulate JAK3-STAT3 and COX-2 pathways. J ETHNOPHARMACOL. 2023 Sep;:117133 IHC,WB;Rat. 37690476