bs-17400R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

HSD17B11 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 51170 SWISS: Q8NBQ5

Target: HSD17B11

Immunogen: KLH conjugated synthetic peptide derived from human Hsd17b11:

201-300/300.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Short-chain alcohol dehydrogenases, such as HSD17B11,

metabolize secondary alcohols and ketones (Brereton et al., 2001

[PubMed 11165019]).[supplied by OMIM, Jun 2009]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Rat (predicted: Human,


Mouse, Rabbit, Pig, Sheep,

Cow, Horse)

Predicted 31 kDa

Subcellular Location: Secreted

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HSD17B11) Polyclonal Antibody, Unconjugated (bs-17400R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.