

bs-17402R**[Primary Antibody]****HSD17B13 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 345275**SWISS:** Q7Z5P4**Target:** HSD17B13**Immunogen:** KLH conjugated synthetic peptide derived from human HSD17B13: 171-270/300.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: 17 beta-HSD13 (17 beta hydroxysteroid dehydrogenase type 13), also designated Short-chain dehydrogenase/reductase 9 (SCDR9), belongs to the 17 beta-HSD family of proteins, which regulate the availability of steroids within various tissues throughout the body. 17 beta-HSD13 is a 300 amino acid secreted protein that is highly expressed in liver and is also detected in ovary, bone marrow, kidney, brain, lung, skeletal muscle, bladder and testis. The gene encoding 17 beta-HSD13 maps to chromosome 4, which houses nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes. Defects in some of the genes located on chromosome 4 are associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.**Applications:** **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Mouse, Rat
(predicted: Human, Rabbit, Pig, Dog, Horse)**Predicted
MW.:** 34 kDa**Subcellular
Location:** Secreted**— VALIDATION IMAGES —**

Sample: Pancreas (Mouse) Lysate at 40 ug
 Primary: Anti-HSD17B13 (bs-17402R) at 1/1000
 dilution Secondary: IRDye800CW Goat Anti-
 Rabbit IgG at 1/20000 dilution Predicted band
 size: 34 kD Observed band size: 34 kD



Sample: Lane 1: Mouse Kidney tissue lysates
 Lane 2: Rat Liver tissue lysates Lane 3: Rat
 Kidney tissue lysates Primary: Anti-HSD17B13
 (bs-17402R) at 1/1000 dilution Secondary:
 IRDye800CW Goat Anti-Rabbit IgG at 1/20000
 dilution Predicted band size: 34 kDa Observed
 band size: 31 kDa



Paraformaldehyde-fixed, paraffin embedded
 (Mouse testis); Antigen retrieval by boiling in
 sodium citrate buffer (pH6.0) for 15min; Block
 endogenous peroxidase by 3% hydrogen
 peroxide for 20 minutes; Blocking buffer (normal
 goat serum) at 37°C for 30min; Antibody
 incubation with (HSD17B13) Polyclonal
 Antibody, Unconjugated (bs-17402R) at 1:400
 overnight at 4°C, followed by operating
 according to SP Kit(Rabbit) (sp-0023)
 instructions and DAB staining.