## bs-5985R

# [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# ABI3 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 51225 SWISS: Q9P2A4

Target: ABI3

**Immunogen:** KLH conjugated synthetic peptide derived from human ABI3:

151-250/366.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: ABI3 is thought to inhibit tumor metastasis. In vitro, ABI3 reduces

cell motility.

Applications: WB (1:500-2000)

**IHC-P** (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Human, Mouse


(predicted: Rat, Rabbit, Pig,

Sheep, Cow, Dog)


Predicted MW.: 39 kDa

Subcellular Cytoplasm Location:


### - VALIDATION IMAGES -



25 ug total protein per lane of various lysates (see on figure) probed with ABI3 polyclonal antibody, unconjugated (bs-5985R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



25 ug total protein per lane of various lysates (see on figure) probed with ABI3 polyclonal antibody, unconjugated (bs-5985R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-ABI3 Polyclonal Antibody, Unconjugated(bs-5985R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining