

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

GDF7 Rabbit pAb

Catalog Number: bs-11461R

Target Protein: GDF7
Concentration: 1mg/ml

Form: Liquid
Host: Rabbit
Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:100-500),

ELISA (1:5000-10000)

Reactivity: (predicted:Human, Mouse, Rat, Pig, Cow, Chicken, Horse)

Predicted MW: 14 kDa Entrez Gene: 151449 Swiss Prot: Q7Z4P5

Source: KLH conjugated synthetic peptide derived from human GDF7: 362-410/450.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Growth/differentiation factors (GDFs) are members of the TGF superfamily (1,2). Members of

the TGF superfamily are involved in embryonic development and adult tissue homeostasis (1). GDF-1 expression is almost exclusively restricted to the central nervous system and mediates cell differentiation events during embryonic development (3). Neither GDF-3 (Vgr-2) nor GDF-9 contains the conserved cysteine residue which is found in most other TGF superfamily members. GDF-3 is detectable in bone marrow, spleen, thymus and adipose tissue, whereas GDF-9 has only been detected in ovary (4). GDF-5 (also designated CDMP-1) has been shown to induce activation of plasminogen activator, thereby inducing angiogenesis. It is predominantly expressed in long bones during fetal embryonic development and is involved in bone formation. (5). GDF-5 mutations have been identified in mice with the mutation brachypodism (bp), a mutation which affects the length and

number of bones in limbs (6). GDF-6 and GDF-7 are closely related to GDF-5 (6). GDF-8 has

been shown to be a negative regulator of skeletal muscle mass (1).

PRODUCT SPECIFIC PUBLICATIONS

| [IF=4.522] Wang Y et al. Aspirin promotes tenogenic differentiation of tendon stem cells and facilitates tendinopathy healing through |
|---|
| regulating the GDF7/Smad1/5 signaling pathway. J Cell Physiol. 2019 Oct 21. WB; Rat . 31637734 |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |