

bs-11469R**[Primary Antibody]****Oligophrenin 1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 4983	SWISS: O60890	
Target: Oligophrenin 1		Reactivity: Rat (predicted: Human, Mouse, Sheep, Cow, Dog, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human Oligophrenin 1: 418-465/802.		
Purification: affinity purified by Protein A		Predicted MW.: 92 kDa
Concentration: 1mg/ml		Subcellular Location: Cell membrane ,Cytoplasm
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Ras p21 can exist in either a physiologically quiescent GDP-binding state or a GTP-binding signal-emitting state (1,2). Interaction of Ras p21 with GTPase activating protein (GAP) can increase the rate of hydrolysis of Ras p21-bound GTP by as much as 1000-fold (3). In mitogenically activated and tyrosine kinase-transformed cells, Ras GAP forms a complex with a protein designated p190 (4). At its amino terminus, p190 contains sequence motifs characteristic of all known GTPases, whereas the carboxy terminus contains sequences similar to those found in the Bcr gene product, n-chimerin and Rho GAP, all of which exhibit intrinsic GAP activity (4,5). Oligophrenin-1 is an additional protein with GTPase activating activity. Oligophrenin-1 is a RhoGAP protein that stimulates GTP hydrolysis of Rho subfamily members and is involved in cell migration, morphogenesis and axon outgrowth (6).		

— VALIDATION IMAGES —

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Oligophrenin 1 Polyclonal Antibody, Unconjugated(bs-11469R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining