

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

CDKN1B/p27 KIP 1 Mouse mAb

Catalog Number: bsm-33425M

Target Protein: CDKN1B/p27 KIP 1

Concentration: 1mg/ml

Form: Size:50ul/100ul/200ul

Liquid

Size: 200ug (PBS only)

Lyophilized

Note: Centrifuge tubes before opening. Reconstitute the lyophilized product in distilled

water. Optimal concentration should be determined by the end user.

Host: Mouse

Clonality: Monoclonal

Clone No.: 1A7
Isotype: IgG1

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human
Predicted MW: 22 kDa
Entrez Gene: 1027
Swiss Prot: P46527

Purification: affinity purified by Protein A

Storage: Size:50ul/100ul/200ul

 $0.01\mbox{M}$ TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Size: 200ug (PBS only)

0.01M PBS


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Cell cycle progression is regulated by cyclins and their cognate Cdks. p27 KIP 1 is a cell cycle

regulatory mitotic inhibitor of cdk activity. p27 KIP 1 is a candidate tumor suppressor gene, and has been proposed to function as a possible mediator of TGF beta induced G1 arrest. p27 KIP 1 is up regulated in response to antimitogenic stimuli. The increased protein expression of p27 results in cellular arrest by binding to cyclin/Cdk complexes such as cyclin

D1/Cdk4. p27 Kip1 is regulated by phosphorylation on serine 10 (S10) and threonine 187 (T187). Phosphorylation by CDK2 on T187 results in ubiquitylation and degradation of p27 Kip 1; while phosphorylation by hKIS on S10 signals the nuclear export to the cytoplasm.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (CDKN1B/p27 KIP 1) Monoclonal Antibody, Unconjugated (bsm-33425M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.