

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **HER2 Rabbit pAb**

Catalog Number: bs-2048R

Target Protein: HER2
Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: ICC/IF (1:100)

Reactivity: Human (predicted:Mouse, Rat, Dog)

Predicted MW: 138 kDa

Entrez Gene: 2064

Swiss Prot: P04626

Source: KLH conjugated synthetic peptide derived from human HER2 receptor: 331-430/1255.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a member of the epidermal growth factor (EGF) receptor family of

receptor tyrosine kinases. This protein has no ligand binding domain of its own and

therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound

EGF receptor family members to form a heterodimer, stabilizing ligand binding and

enhancing kinase-mediated activation of downstream signalling pathways, such as those

involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic


variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform

b) have been reported, with the most common allele, Ile654/Ile655, shown here.

Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully

characterized. [provided by RefSeq, Jul 2008].

## **VALIDATION IMAGES**



Tissue/cell:A549 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Antibody incubation with (HER2 receptor) polyclonal Antibody, Unconjugated (bs-2048R) 1:100, 90 minutes at 37°C; followed by a FITC conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.