bsm-52253R

[Primary Antibody]

NGFR/p75NTR Recombinant Rabbit mAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 5D4
GeneID: 4804 SWISS: P08138

Target: NGFR/p75NTR

Immunogen: A synthesized peptide derived from human p75 NTR: 350-400.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

induced apoptosis.

Background: The low affinity NGFR (Nerve growth factor receptor) is a 75kDa

membrane-spanning glycoprotein lacking intrinsic tyrosine kinase activity. p75NGFR interacts with TrkA, the high affinity NGF receptor and potentiates TrkA signaling at low NGF concentrations. The p75 receptor binds nerve growth factor, brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4 with varying specificities. The p75NGFR plays an important role in neurotrophic factor signaling and has been shown to modulate the susceptibility of selective cellular populations to programmed cell death. It is expressed on many neuronal cells types including many embryonic forms and the receptor can be used to isolate neuronal progenitor cells. NGF is important for the development, differentiation and survival of a variety of neuronal and nonneuronal cells. Its action is mediated by binding to two distinct receptors, the high affinity p140 and low affinity p75. p75NGFR binds neurotrophins including brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3), NT-4/5, and NT-6. p75NGFR belongs to the TNF-R superfamily and is reported to mediate NGF-

Applications: IHC-P (1:100-500)


IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:50-200)

Reactivity: Human, Mouse, Rat

Predicted MW.: 44 kDa


Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES




Paraformaldehyde-fixed, paraffin embedded Human Prostate; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with NGFR/p75NTR Monoclonal Antibody,

Unconjugated(bsm-52253R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.





Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with NGFR/p75NTR Monoclonal Antibody,

Unconjugated(bsm-52253R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



4% Paraformaldehyde-fixed PC-12 (Rat) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (NGFR/p75NTR) monoclonal Antibody, unconjugated (bsm-52253R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green,bs-40295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.





Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NGFR/p75NTR Monoclonal Antibody, Unconjugated (bsm-52253R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Red, bs-0295G-BF594), DAPI (blue, C02-04002) was used to stain the cell nuclei.

Paraformaldehyde-fixed, paraffin embedded Mouse Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NGFR/p75NTR Monoclonal Antibody, Unconjugated (bsm-52253R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Red, bs-0295G-BF594), DAPI (blue, C02-04002) was used to stain the cell nuclei.