## bs-9860R

## [ Primary Antibody ]



## phospho-cardiac Troponin I (Thr143) Rabbit pAb A N T B =

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

**Host:** Rabbit **Isotype:** IgG

Clonality: Polyclonal

**GenelD:** 7137 **SWISS:** P19429

Target: cardiac Troponin I (Thr143)

Immunogen: KLH conjugated synthesised phosphopeptide derived from human

cardiac Troponin I around the phosphorylation site of Thr143:

RP(p-T)LR.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Troponin I (TnI), along with troponin T (TnT) and troponin C (TnC), is one of 3 subunits that form the troponin complex of the thin

filaments of striated muscle. TnI is the inhibitory subunit; blocking actin-myosin interactions and thereby mediating striated muscle relaxation. The TnI subfamily contains three genes: TnI-skeletal-fast-twitch, TnI-skeletal-slow-twitch, and TnI-cardiac. This gene encodes the TnI-cardiac protein and is exclusively expressed in cardiac muscle tissues. Mutations in this gene cause familial hypertrophic cardiomyopathy type 7 (CMH7) and familial

restrictive cardiomyopathy (RCM). [provided by RefSeq, Jul 2008]

**Applications: IHC-P** (1:100-500)

IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test)

Reactivity: Human, Rat

(predicted: Mouse, Pig, Sheep, Cow, Dog, Horse)

Predicted 24


**MW.:** <sup>24 kDa</sup>

Subcellular Cytoplasm Location:


## VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PGGT1B) Polyclonal Antibody, Unconjugated (bs-9860R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-cardiac Troponin I (Thr143)) Polyclonal Antibody, Unconjugated (bs-9860R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Blank control: K562. Primary Antibody (green line): Rabbit Anti-phospho-cardiac Troponin I (Thr143) antibody (bs-9860R) Dilution: 1µg/10^6 cells: Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat antirabbit IgG-FITC Dilution:  $1\mu g$  /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block nonspecific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.