

bs-9865R**[Primary Antibody]****MYLK3 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 91807**SWISS:** Q32MK0**Target:** MYLK3**Immunogen:** KLH conjugated synthetic peptide derived from human MYLK3: 701-800/817.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The Ca²⁺/calmodulin-dependent protein kinases (CaM kinases) are a structurally related subfamily of serine/threonine kinases that includes CaMKI, CaMKII, CaMKIV and Myosin light chain kinases (MYLKs, also designated MLCKs). The MYLK kinases phosphorylate Myosin regulatory light chains to catalyze Myosin interaction with Actin filaments, resulting in contractile activity. MYLK3 (Myosin light chain kinase 3) is a 795 amino acid cardiac-specific protein that contains one protein kinase domain and belongs to the protein kinase superfamily. Like other MYLK kinases, MYLK3 is thought to play a role in smooth muscle contraction, specifically using magnesium as a cofactor to catalyze the ATP-dependent phosphorylation of Myosin light chain proteins. Additionally, MYLK3 may regulate sarcomere assembly in heart tissue, possibly mediating proper heart function.

Applications: WB (1:500-2000)**Reactivity:** Mouse (predicted: Human, Rat, Rabbit, Sheep, Cow, Dog, Horse)**Predicted MW.:** 98 kDa**Subcellular Location:** Cytoplasm**— VALIDATION IMAGES —**

25 ug total protein per lane of various lysates (see on figure) probed with MYLK3 polyclonal antibody, unconjugated (bs-9865R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Sample: Bone (Mouse) Lysate at 40 ug Primary: Anti- MYLK3 (bs-9865R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 98 kD
Observed band size: 100 kD