


bs-41495R**[Primary Antibody]****THRB Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 7068**SWISS:** P10828**Target:** THRB**Immunogen:** Recombinant human THRB protein : 209-461/461.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** Thyroid hormone receptors (TRs) are ligand-dependent transcription factors that mediate the biological activities of thyroid hormone (T3). Thyroid hormone receptor b2 (TRb2) is a high affinity receptor for triiodothyronine which belongs to the nuclear hormone receptor family and the NR1 subfamily. It is composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal steroid-binding domain. Defects in the receptor result in generalized thyroid hormone resistance (GTHR). GTHR is transmitted as an autosomal dominant trait, but an autosomal recessive form also exists. The disease is characterized by goiter, abnormal mental functions, increased susceptibility to infections, abnormal growth and bone maturation, tachycardia and deafness. GTHR patients also have high levels of circulating thyroid hormones (T3-T4), with normal or slightly elevated thyroid stimulating hormone.**Applications:** IHC-P (1:200-800)**IHC-F** (1:400-800)**IF** (1:100-500)**Flow-Cyt** (1ug/Test)**ELISA** (1:5000-10000)**Reactivity:** Human**Predicted MW.:** 53 kDa**Subcellular Location:** Nucleus**— VALIDATION IMAGES —**


Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (THRB) Polyclonal Antibody, Unconjugated (bs-41495R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human thyroid gland); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (THRB) Polyclonal Antibody, Unconjugated (bs-41495R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control (black line) :Hela. Primary Antibody (green line): Rabbit Anti-THRB antibody (bs-41495R) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control (black line) :Hela. Primary
 Antibody (green line): Rabbit Anti-THRB
 antibody (bs-41495R) Dilution:1ug/Test;
 Secondary Antibody (white blue line) : Goat
 anti-rabbit IgG-AF488 Dilution: 0.5ug/Test.
 Isotype control (orange line) : Normal Rabbit
 IgG Protocol The cells were fixed with 4% PFA
 (10min at room temperature)and then
 permeabilized with 90% ice-cold methanol for
 20 min at -20°C, The cells were then incubated in
 5%BSA to block non-specific protein-protein
 interactions for 30 min at room temperature
 .Cells stained with Primary Antibody for 30 min
 at room temperature. The secondary antibody
 used for 40 min at room temperature.
 Acquisition of 20,000 events was performed.