


bs-2301R**[Primary Antibody]****CD162 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 6404**SWISS:** Q14242**Target:** CD162**Immunogen:** KLH conjugated synthetic peptide derived from human PSGL-1: 251-350/412. < Extracellular >**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** This gene encodes a glycoprotein that functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L-selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking during inflammation by tethering of leukocytes to activated platelets or endothelia expressing selectins. This protein requires two post-translational modifications, tyrosine sulfation and the addition of the sialyl Lewis x tetrasaccharide (sLex) to its O-linked glycans, for its high-affinity binding activity. Aberrant expression of this gene and polymorphisms in this gene are associated with defects in the innate and adaptive immune response. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Apr 2011].**Applications:** WB (1:500-2000)**Flow-Cyt** (1µg /test)**ICC/IF** (1:25)**Reactivity:** Human**Predicted MW.:** 41 kDa**Subcellular Location:** Cell membrane**— VALIDATION IMAGES —**


Sample: Jurkat(Human) Cell Lysate at 30 ug
MOLT-4(Human) Cell Lysate at 30 ug Primary:
Anti-CD162 (bs-2301R) at 1/500 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 41 kD
Observed band size: 41 kD



Sample: Raji(Human) Cell Lysate at 30 ug
U937(Human) Cell Lysate at 30 ug Primary: Anti-
CD162 (bs-2301R) at 1/500 dilution Secondary:
IRDye800CW Goat Anti-Rabbit IgG at 1/20000
dilution Predicted band size: 41 kD Observed
band size: 41 kD



MCF7 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (CD162) polyclonal Antibody, Unconjugated (bs-2301R) 1:25, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control: U937(blue). Primary Antibody:
Rabbit Anti-CD162 antibody(bs-2301R), Dilution:
1µg in 100 µL 1X PBS containing 0.5% BSA;

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Isotype Control Antibody: Rabbit IgG (orange)
,used under the same conditions. Secondary
Antibody: Goat anti-rabbit IgG-PE(white blue),
Dilution: 1:200 in 1 X PBS containing 0.5% BSA.
Protocol The cells were fixed with 2%
paraformaldehyde (10 min).Primary antibody
(bs-2301R, 1µg /1x10⁶ cells) were incubated for
30 min on the ice, followed by 1 X PBS
containing 0.5% BSA + 10% goat serum (15 min)
to block non-specific protein-protein
interactions. Then the Goat Anti-rabbit IgG/PE
antibody was added into the blocking buffer
mentioned above to react with the primary
antibody at 1/200 dilution for 30 min on ice.
Acquisition of 20,000 events was performed.